

Filed: 31-Dec-2001

In the United States Patent and Trademark Office

PATENT APPLICATION

TITLE:

METHOD FOR TREATING GLAUCOMA II B

INVENTOR(s):

Dilip Wagle
Martin Gall
Stanley C. Bell
Edmond J. LaVoie

METHOD FOR TREATING GLAUCOMA IIB

The present application claims the priority of US Applications 60/296,258, filed 6
5 June 2001, and 60/259,428, filed 29 December 2000.

The present invention relates to methods for treating glaucoma or improving
accommodation (i.e. the process by which the eye adjusts for vision at different
distances), and to compounds and compositions for use in such treating. In one aspect,
the present invention relates to a method of decreasing the intraocular pressure caused by
10 glaucoma.

Diabetes is the major determinant to the development of visual disability and
blindness in parts of the world unencumbered by causes related to malnutrition or
infectious diseases. Retinopathy is the leading cause of blindness in diabetics and is a
progressive, degenerative disease. Of the many risk factors believed to be associated
15 with diabetic retinopathy, the level of glucose in the plasma has been widely
investigated. It is well accepted that a lower incidence of retinopathy is associated with
decreased plasma levels of glucose.

Ophthalmologic disorders in diabetes include opacification and glaucoma. As the
occurrence of these indications is correlated with the persistent hyperglycemia of the
20 disease. Although the incidence of glaucoma is significant in diabetic populations,
glaucoma affects a substantial portion of the general aging population as well.

Primary open angle glaucoma occurs in approximately 4% of diabetics compared
to 1.8% of the general population. The reasons for the increase in intraocular pressure
that is observed in this disorder are not completely understood. The increase in
25 intraocular pressure that characterizes glaucoma is likely caused by an impairment in the
drainage of fluid from the eye at the trabecular meshwork since trabeculectomy restores,
at least for a period of time, normal intraocular pressures. The origin of this impairment
to fluid movement is currently unknown but may be related to a physical obstruction or
restriction to movement of proteins that make up a sieving system in the trabecular
30 meshwork. The trabecular meshwork functions as a sieving system that maintains a
restricted flow of intraocular fluid from the eye. The result of excess restriction of this
flow is a back pressure that causes increased intraocular pressure.


Replacement of the trabecular meshwork (trabeculectomy) remains an established surgical procedure for improving the filtering of intraocular fluid and for overall reduction of intraocular pressure. This remedy is invasive and of limited effectiveness, since pressure elevation frequently recurs after the procedures.

5 Current chronic pharmaceutical therapies impose a measure of risk on an already medically compromised patient population. The use of topical B-blockers may affect underlying cardiovascular disease, and carbonic anhydrase inhibitors (e.g. Diamox™) may cause metabolic acidosis. The use of pressure-lowering drugs will be affected by the state of renal disease in compromised elderly and diabetic patients. The drawbacks
10 associated with current pharmaceutical therapies highlight an unmet medical need for a chronic pharmaceutical intervention that is distinct in mechanism of action from current therapies.

New strategies for pharmaceutical intervention in the treatment of glaucoma based upon new mechanisms of action need to be identified. In addition, pharmaceutical
15 agents that decrease the intraocular pressure associated with glaucoma are needed. Also, the methods of improving accommodation provided by the invention allow one to avoid costly and burdensome optical solutions, such as the use of separate reading glasses or glasses with bifocal lenses.

Summary of the Invention

20 In one embodiment, the invention relates to a method of treating or ameliorating or preventing glaucoma, decreasing intraocular pressure or improving or ameliorating ocular accommodation in an animal, including a human, comprising administering an intraocular pressure decreasing amount or ocular accommodation improving amount of a compound of the formula I or IA,



wherein the substituent groups are as defined below.

Detailed Description of the Invention

In accordance with the present invention a method is provided for the treatment of an animal, preferably a mammal, preferably a human with ophthalmologic disorders including glaucoma and reduced accommodation. Briefly the method of the present invention provides for a method of treatment of mammals with glaucoma or reduced accommodation that can be caused by age or certain age-related diseased states such as diabetes. The method provides for administration of classes of inhibitors of advanced glycation. The invention further provides for methods to monitor the improvement in the ocular condition during the course of the administration of compound.

10 The agents used in the invention are compounds of formulas I or IA, wherein:

- a. J is oxygen, sulfur, or N-R^d;
- b. the carbon 2 to nitrogen bond is a double bond except when R^c is oxo;
- c. the bond between carbons 4 and 5 is a single bond or a double bond (in one embodiment, a single bond);

15 d. R^a and R^b are

- 1. independently selected from hydrogen, acylamino, acyloxyalkyl, alkanoyl, alkanoylalkyl, alkenyl, alkoxy, alkoxy carbonyl, alkoxy carbonylalkyl, alkyl, alkylamino, (C₁-C₃)alkylenedioxy, allyl, amino, ω-alkylenesulfonic acid, carbamoyl, carboxy, carboxyalkyl (which alkyl can be substituted with alkyloxyimino), cycloalkyl, dialkylamino, halo, hydroxy, (C₂-C₆)hydroxyalkyl, mercapto, nitro, sulfamoyl, sulfonic acid, alkylsulfonyl, alkylsulfinyl, alkylthio, trifluoromethyl, morpholin-4-yl, thiomorpholin-4-yl, piperidin-1-yl, 4-[C₆ or C₁₀] arylpiperidin-1-yl, 4-[C₆ or C₁₀] arylpiperazin-1-yl, Ar {wherein, consistent with the rules of aromaticity, Ar is C₆ or C₁₀ aryl or a 5- or 6-membered heteroaryl ring, wherein the 6-membered heteroaryl ring contains one to three atoms of N, and the 5-membered heteroaryl ring contains from one to three atoms of N or one atom of O or S and zero to two atoms of N, each heteroaryl ring can be fused to a substituted benzene, pyridine, pyrimidine, pyrazine, pyridazine, or (1,2,3)triazine (wherein the ring fusion is at a carbon-carbon double bond of Ar)(or wherein Ar is as above but not heteroaryl fused to pyridine)} (in one embodiment, Ar is C₆ or C₁₀ aryl)}, Ar-alkyl, ArO-, ArSO₂-, ArSO-, ArS-, ArSO₂NH-, ArNH-, (N-Ar)(N-alkyl)N-, ArC(O)-, ArC(O)NH-, ArNH-C(O)-, and (N-Ar)(N-alkyl)N-C(O)-, or

together R^a and R^b comprise methylenedioxy- (in one embodiment, R^a and R^b are not acyloxyalkyl, alkenyl, (C₁-C₃)alkylenedioxy or allyl, or together do not comprise methylenedioxy); or

2. together with their ring carbons form a C₆- or C₁₀- aryl fused ring; or

5 3. together with their ring carbons form a C₅-C₇ fused cycloalkyl ring having up to two double bonds including any fused double bond of the containing group, which cycloalkyl ring can be substituted by one or more of the group consisting of alkyl, alkoxy carbonyl, amino, aminocarbonyl, carboxy, fluoro, or oxo (in one embodiment, the ring having no double bonds except any fused double bond of the formula I or IA ring, which cycloalkyl ring can be substituted by one or more of the group consisting of alkyl, amino, aminocarbonyl, carboxy, fluoro, or oxo, where multiple substituents are located on different carbon atoms of the cycloalkyl ring, except in the case of alkyl and fluoro substituents, which can be located on the same or different carbon atoms); or

15 4. together with their ring carbons form a fused 5- or 6-membered heteroaryl ring, wherein the 6-membered heteroaryl ring contains one to three atoms of N, and the 5-membered heteroaryl ring contains from one to three atoms of N or one atom of O or S and zero to two atoms of N; or

20 5. together with their ring carbons form a fused five to eight membered second heterocycle (in one embodiment, a fused five to six membered second heterocycle), wherein the fused heterocycle consists of ring atoms selected from the group consisting of carbon, nitrogen, oxygen, sulfur, and S(O)_n, wherein n is 1 or 2;

e. R^d is alkyl, alkenyl, hydrogen, or Ar;

25 f. R^c is

1. oxo (when $\Delta^{2,3}$ is not present), or (when $\Delta^{2,3}$ is present) hydrogen, alkyl, alkylthio, hydrogen, mercapto, amino, amino(C₁-C₅)alkyl, or amino(C₆ or C₁₀)aryl, or wherein the amino of the last three groups can be substituted with

(a) Ar,

30 (b) Ar-Z-, Ar-alkyl-Z-, Ar-Z-alkyl, Ar-amino-Z-, Ar-aminoalkyl-Z-, or Ar-oxyalkyl-Z-, wherein Z is a carbonyl or -SO₂-


(c) formyl or alkanoyl (in one embodiment, R^c is according to one of these optionally substituted three groups; in another, J is S or O (in one embodiment S), and R^c is hydrogen, oxo, alkyl, amino, amino(C_1-C_5)alkyl or aminophenyl, wherein the amino of the latter three groups can be substituted as outlined here),

2. $-NHC(O)(CH_2)_n-D-R^eR^f$, wherein D is oxygen, sulfur or nitrogen, wherein where D is nitrogen n is 0, 1 or 2, but when D is oxygen or sulfur n=1 or 2, and R^f is present only when D is nitrogen, wherein

(a) R^e is

(1) Ar,

(2) a group of the formula



wherein E is sulfur, oxygen, or $N-R^i$, and R^g , R^h and R^i are independently the same as R^a , R^b and R^d , respectively,

(3) a C_3-C_8 cycloalkyl ring having up to one double bond with the proviso that the carbon linking the cycloalkyl ring to D is saturated, which cycloalkyl ring can be substituted by one or more alkyl-, alkoxy-, amino-, aminocarbonyl-, carboxy-, fluoro-, or oxo-substituents;

(4) a 5- or 6-membered heteroaryl ring containing at least one and up to three atoms of N for the 6-membered heteroaryl rings and from one to three atoms of N or one atom of O or S and zero to two atoms of N for the 5-membered heteroaryl rings;

(5) hydrogen, (C_2-C_6)hydroxyalkyl, alkanoylalkyl, alkyl, alkoxyalkyl, alkenyl, carboxyalkyl (which alkyl can be substituted with alkoxyimino), alkoxyalkyl, a group Ar^ϕ which is C_6 - or C_{10} - aryl or a 5- or 6-membered, or 9- or 10-membered

heteroaryl (wherein the heteroatom is one oxygen, one sulfur or one nitrogen) or Ar^ϕ -alkyl; and

(b) R^f is independently hydrogen, $(\text{C}_2\text{-C}_6)$ hydroxyalkyl, alkanoylalkyl, alkyl, alkoxyalkyl, alkenyl, carboxyalkyl (which alkyl can be substituted with alkoxyimino), alkoxyalkyl, Ar^ϕ , or Ar^ϕ -alkyl;

wherein aryl, Ar, or Ar^ϕ can be substituted with, in addition to any substitutions specifically noted one or more general substituents selected from the group of acylamino, acyloxyalkyl, alkanoyl, alkanoylalkyl, alkenyl, alkoxy, alkoxyalkyl, alkoxyalkylalkyl, alkyl, alkylamino, $(\text{C}_1\text{-C}_3)$ alkylenedioxy, alkylsulfonyl, alkylsulfinyl, ω -alkylenesulfonic acid, alkylthio, allyl, amino, $\text{ArC}(\text{O})$ -, $\text{ArC}(\text{O})\text{NH}$ -, carboxy, carboxyalkyl, cycloalkyl, dialkylamino, halo, trifluoromethyl, hydroxy, $(\text{C}_2\text{-C}_6)$ hydroxyalkyl, mercapto, nitro, ArO -, Ar-, Ar-alkyl-, sulfamoyl, sulfonic acid, 1-pyrrolidinyl, 4- $[\text{C}_6$ or $\text{C}_{10}]$ aryl piperazin-1-yl, 4- $[\text{C}_6$ or $\text{C}_{10}]$ aryl piperidin-1-yl, azetidin-1-yl, morpholin-4-yl, thiomorpholin-4-yl, piperidin-1-yl (in one embodiment, the substituents are selected from the group of aryl preferred general substituents which are: alkyl, amino, dialkylamino, 1-pyrrolidinyl, 4- $[\text{C}_6$ or $\text{C}_{10}]$ aryl piperazin-1-yl, 4- $[\text{C}_6$ or $\text{C}_{10}]$ aryl piperidin-1-yl, azetidin-1-yl, morpholin-4-yl, thiomorpholin-4-yl and piperidin-1-yl); and

heterocycles, except those of Ar and Ar^ϕ , can be substituted with in addition to any substitutions specifically noted one or more preferred substituents selected from acylamino, alkanoyl, alkoxy, alkoxyalkyl, alkoxyalkylalkyl, alkyl, $(\text{C}_1$ to $\text{C}_3)$ alkylenedioxy, alkylamino, alkylsulfonyl, alkylsulfinyl, alkylthio, amino, $\text{ArC}(\text{O})$ -, ArO -, Ar-, Ar-alkyl, carboxy, dialkylamino, fluoro, fluoroalkyl, difluoroalkyl, hydroxy, mercapto, oxo, sulfamoyl, trifluoromethyl, 4- $[\text{C}_6$ or $\text{C}_{10}]$ aryl piperidin-1-yl and 4- $[\text{C}_6$ or $\text{C}_{10}]$ aryl piperazin-1-yl (in one embodiment, the substituents selected from heterocycle preferred general substituents: acylamino, alkanoyl, alkoxy, alkoxyalkyl, alkoxyalkylalkyl, alkyl, $(\text{C}_1$ to $\text{C}_3)$ alkylenedioxy, alkylamino, alkylsulfonyl, alkylsulfinyl, alkylthio, amino, $\text{ArC}(\text{O})$ -, ArO -, Ar-, Ar-alkyl, carboxy, dialkylamino, fluoro, fluoroalkyl, difluoroalkyl, hydroxy, mercapto, oxo, sulfamoyl, trifluoromethyl, 4- $[\text{C}_6$ or $\text{C}_{10}]$ aryl piperidin-1-yl and 4- $[\text{C}_6$ or $\text{C}_{10}]$ aryl piperazin-1-yl, wherein multiple

substituents are located on different atoms of the heterocyclic ring, with the proviso that alkyl, alkoxycarbonyl, and fluoro substituents can be substituted on the same carbon atom of the heterocyclic ring);

or a pharmaceutically acceptable salt of said compounds,

- 5 with the proviso that where the compound of formula I is administered to decrease intraocular pressure at least one compound of formula I administered in effective amount is not a thiazole substituted on a ring carbon sulfonamide (the amide of which can be substituted) that has carbonic anhydrase inhibiting activity.

- 10 Primary open angle glaucoma is characterized by an increase in intraocular pressure. The condition of open angle glaucoma is characterized by an increase in the pressure within a person's eye or eyes, called the intraocular pressure. The normal pressure is about 15 mmHg. Elevated pressures of 20-30 mm Hg create a strong risk of damage to the optic nerve and blindness.

- 15 Glucose reacts with proteins by a non-enzymatic, post-translational modification process called non-enzymatic glycosylation. The resulting sugar-derived adduct, the advanced glycosylation end product (AGE), matures to a molecular species that is reactive, and can readily bond to amino groups on adjacent proteins, resulting in the formation of AGE cross-links between proteins.

- 20 It has now been found that certain compounds that inhibit the formation of such sugar-derived adducts, or in some cases are believed to deactivate such adducts or break resulting crosslinks, can reduce intraocular pressure or ameliorate a trend towards elevated pressure.

- Structural matrix proteins isolated from tissues of diabetics and aged individuals
25 are more highly crosslinked than those from nondiabetics or younger individuals and are more resistant to both enzymatic and chemical hydrolysis *in vitro*. It is this cross-linked state of proteins that is believed to cause stiffness of tissues. The cleavage of AGE cross-links between proteins can provide a mechanism-based therapy for restoration of normal tissue function. An agent that cleaves AGE cross-links between proteins or
30 inhibits their formation can restore more normal sieving function and movement to the trabecular meshwork.

 In accordance with the present invention, methods for administering pharmaceutical compositions containing compounds have been developed for the

treating glaucoma, intraocular pressure associated with glaucoma, and reduced accommodation. These agents are either substituted thiazole, oxazole, or imidazole agents as shown in the Summary section above.

As is noted in the formula for I and IA, the invention includes aromatic thiazole, oxazole, and imidazole analogs, as well as non aromatic analogs thereof such as thiazoline, thiazolidine, oxazoline, oxazolidine, imidazoline, and imidazolidine analogs.

The alkyl, and alkenyl groups referred to above include both C1 to C6 linear and branched alkyl and alkenyl groups, unless otherwise noted. Alkoxy groups include linear or branched C1 to C6 alkoxy groups, unless otherwise noted.

“Ar” (consistent with the rules governing aromaticity) refers to a C₆ or C₁₀ aryl, or a 5 or 6 membered heteroaryl ring. The heteroaryl ring contains at least one and up to three atoms of N for the 6 membered heteroaryl ring. The 5 membered heteroaryl ring contains; (1) from one to three atoms of N, or (2) one atom of O or S and zero to two atoms of N. Nonlimiting examples of heteroaryl groups include: pyrrolyl, furanyl, thienyl, pyridyl, oxazolyl, pyrazolyl, pyrimidinyl, and pyridazinyl.

“Ar” can be fused to either a benzene, pyridine, pyrimidine, pyridazine, or (1,2,3) triazine ring.

As used herein, C₆ or C₁₀ aryl groups and heteroaryl containing five or six, or nine to ten ring members are monocyclic or bicyclic.

In certain embodiments of the invention, the thiazoles, imidazoles, and oxazoles of the invention contain R^a and R^b substitutions that together with their ring carbons (the C4-C5 carbons of the thiazoles, imidazoles, and oxazoles) form a five to eight membered fused heterocycle (i.e. a bicyclic heterocycle is formed). In these embodiments the fused heterocycle is preferably not aromatic. Particular compounds within these embodiments contain sulfur atoms in the fused heterocycle (the ring fused to the thiazoles, imidazoles, and oxazoles). These sulfur atoms in these particular compounds can exist in various oxidation states, as S(O)_n, where n is 0, 1, or 2.

In certain embodiments of the invention, thiazoles, imidazoles, and oxazoles of the invention contain R^a and R^b substitutions that together with their ring carbons (the C4-C5 carbons of the thiazoles, imidazoles, and oxazoles) form a C5 to C7 cycloalkyl ring having up to double bonds including the C4-C5 double bond. In other embodiments a cycloalkyl ring is present when R^c is a C3 to C8 cycloalkyl ring. The cycloalkyl groups can be substituted by one or more of the group consisting of alkyl-,

alkoxycarbonyl-, amino-, aminocarbonyl-, carboxy-, fluoro-, or oxo- substituents. One of ordinary skill in the art will recognize that where cycloalkyl groups contain double bonds, the sp^2 hybridized carbon atoms can contain only one substituent (which cannot be amino- or oxo-). Sp^3 hybridized carbon atoms in the cycloalkyl ring can be geminally substituted with the exception that (1) two amino groups and (2) one amino and one fluoro group can not be substituted on the same sp^3 hybridized carbon atom.

In certain embodiments of the invention, the thiazoles, imidazoles, and oxazoles of the invention contain R^a and R^b substitutions that together with their ring carbons (the C4-C5 carbons of the thiazoles, imidazoles, and oxazoles) form a five or six membered heteroaryl ring (i.e, a bicyclic aromatic heterocycle is formed). A preferred bicyclic aromatic heterocycle of the invention is a purine analog [J is N- R^d and R^a and R^b together with their ring carbons (the C4 and C5 of the imidazole ring) form a pyrimidine ring].

Aryl, Ar, or Ar^ϕ can be substituted with, in addition to any substitutions specifically noted one or more substituents selected from the group of acylamino, acyloxyalkyl, alkanoyl, alkanoylalkyl, alkenyl, alkoxy, alkoxycarbonyl, alkoxycarbonylalkyl, alkyl, alkylamino, (C1-C3)alkylenedioxy, alkylsulfonyl [alkylSO₂-], alkylsulfinyl [alkylSO-], ω -alkylenesulfonic acid [-alkylSO₃H where n = 1 to 6], alkylthio, allyl, amino, ArC(O)-, ArC(O)NH-, carboxy, carboxyalkyl, cycloalkyl, dialkylamino, halo, trifluoromethyl, hydroxy, (C2-C6)hydroxyalkyl, mercapto, nitro, ArO-, Ar-, Ar-alkyl-, sulfamoyl, sulfonic acid, morpholin-4-yl, thiomorpholin-4-yl, piperidin-1-yl, 1-pyrrolidinyl, 4-[C₆ or C₁₀]arylpiperidin-1-yl and 4-[C₆ or C₁₀]arylpiperazin-1-yl.

Heterocycles, except those of Ar and Ar^ϕ , can be substituted with in addition to any substitutions specifically noted one or more substituents selected from acylamino, alkanoyl, alkoxy, alkoxycarbonyl, alkoxycarbonylalkyl, alkyl, (C1 to C3)alkylenedioxy, alkylamino, alkylsulfonyl, alkylsulfinyl, alkylthio, amino, ArC(O)-, ArO-, Ar-, Ar-alkyl, carboxy, dialkylamino, fluoro, fluoroalkyl, difluoroalkyl, hydroxy, mercapto, oxo, sulfamoyl, trifluoromethyl, 4-[C₆ or C₁₀]arylpiperidin-1-yl and 4-[C₆ or C₁₀]arylpiperazin-1-yl, wherein multiple substituents are located on different atoms of the heterocyclic ring, with the proviso that alkyl, alkoxycarbonyl, and fluoro substituents

can be substituted on the same carbon atom of the heterocyclic ring. Heterocycles can be substituted with one or more substituents.

The halo atoms can be fluoro, chloro, bromo or iodo. Chloro and fluoro are preferred for aryl substitutions.

5 In some embodiments of this invention, the compounds of formula (I) can form biologically and pharmaceutically acceptable salts. Useful salt forms include the halides (particularly bromides and chlorides), tosylates, methanesulfonates, brosylates, fumarates, maleates, succinates, acetates, mesitylenesulfonates, and the like. Other related salts can be formed using similarly non-toxic, biologically or pharmaceutically
10 acceptable anions.

Representative, non-limiting examples of compounds of the present invention are:

Thiazole

4,5-Dimethylthiazole

15 4-Methylthiazole

5-Methylthiazole

4-Methyl-5-(2-hydroxyethyl)thiazole

4-Methyl-5-vinylthiazole

Benzothiazole

20 2-Aminobenzothiazole

2-Amino-4-chlorobenzothiazole

2-Amino-6-chlorobenzothiazole

2,6-Diamino-benzothiazole

2-Aminothiazole

25 2,4,5-Trimethylthiazole

2-Amino-5-methylthiazole

2-Amino-4-methylthiazole

2-Acetylthiazole

2-Ethyl-4-methylthiazole

30 Ethyl 2-(Formylamino)-4-thiazoleacetate

2-(Formylamino)-alpha-(methoxyimino)-4-thiazoleacetic acid

2-Amino-4-phenylthiazole hydrochloride monohydrate

2-Isobutylthiazole

2-Methyl-2-thiazoline

2-Methyl-2-oxazoline

2-Oxazolidone


2-Amino-4-thiazoleacetic acid

5 1-(Thiazolyl)-3-phenyl-urea

1-(Thiazolidinyl)-3-(4-fluorophenyl)-urea

(4-fluorophenyl)thiazolin-2-ylamine

2-(4,6-dimethylpyrimidin-2-ylthio)-N-(1,3-thiazol-2'yl)acetamide, also known as
N-(Thiazolyl)-2-(4,6-dimethyl-pyrimidin-2-yl-thio)-acetamide



10

2-(4-propylphenoxy)-N-(thiazol-2-yl)acetamide

2-furyl-N-[4-(6-methylbenzothiazol-2-yl)phenyl]carboxamide

2-(3,5-Dimethylphenoxy)-N-thiazol-2-yl)acetamide

5,5-Dimethyl-2-(2-naphthylamino)-4,5,6-trihydrobenzothiazol-7-one

15

Imidazole

1-Methylimidazole

1-Ethylimidazole

1-Butylimidazole

1-Vinylimidazole

20

1-Allylimidazole

1-(Trimethylsilyl) imidazole

1-(3-Aminopropyl) imidazole

1-Benzyl imidazole

1-Phenyl imidazole

25

1,5-Dicyclohexyl imidazole

1-(p-Toluenesulfonyl) imidazole

N-Benzoyl-imidazole

4-Methyl-imidazole

4'-(Imidazol-1-yl)-acetophenone
4-(Imidazol-1-yl)-phenol
1-(4-Methoxyphenyl)-1H-imidazole
Methyl-4-(1H-imidazol-1-yl)benzoate
5 1-Methylbenzimidazole

as well as other biologically and pharmaceutically acceptable salts thereof.

Where one or more compounds of formula I are administered to decrease intraocular pressure, at least one compound of formula I administered in effective
10 amount is not a thiazole substituted on a ring carbon sulfonamide (the amide of which can be substituted) that has carbonic anhydrase inhibiting activity. Of course, the composition can include an effective amount of a first agent, as well as a carbonic anhydrase-inhibiting effective amount of another agent, including one of those distinguished above.

15 Compounds of the formula II can be conveniently prepared by chemical syntheses well-known in the art. Certain of the compounds are well-known and readily available from chemical supply houses or can be prepared by synthetic methods specifically published therefor. For instance, 4,5-Dimethylthiazole, 4-Methylthiazole, 5-Methylthiazole, 4-Methyl-5-thiazoleethanol, 4-Methyl-5-vinylthiazole, Benzothiazole, 2-
20 Aminobenzothiazole, 2-Amino-4-chlorobenzothiazole, 2-Amino-6-chlorobenzothiazole, 2-Aminothiazole, 2,4,5-Trimethylthiazole, 2-Amino-5-methylthiazole, 2-Amino-4-methylthiazole, 2-Acetylthiazole, 2-Ethyl-4-methylthiazole, Ethyl 2-(Formylamino)-4-thiazoleacetate, 2-(Formylamino)-alpha-(methoxyimino)-4-thiazoleacetic acid, 2-amino-4-phenylthiazole hydrochloride monohydrate, 2-
25 Isobutylthiazole, 2-Methyl-2-thiazoline, 2-Methyl-2-oxazoline, 2-Oxazolidone, Thiomorpholine, 2-Amino-4-thiazoleacetic acid, Imidazole, 1-Methylimidazole, 1-Butylimidazole, 1-Vinylimidazole, 1-Allylimidazole, 1-(Trimethylsilyl) imidazole, 1-(3-Aminopropyl) imidazole, 1-Benzyl imidazole, 1-Phenyl imidazole, 1,5-Dicyclohexyl imidazole, 1-(p-Toluenesulfonyl) imidazole, N-Benzoyl-imidazole, 4-Methyl-imidazole,
30 4'-(Imidazol-1-yl)-acetophenone, 4-(Imidazol-1-yl)-phenol, 1-(4-Methoxyphenyl)-1H-imidazol and 1-Methylbenzimidazole can be obtained from Sigma (St. Louis, MO), Aldrich (Milwaukee, WI) or Fluka (Milwaukee, WI) (all divisions of Sigma-Aldrich Co.).

1-ethylimidazole can be obtained from TCI America (Portland, OR). N-(Thiazolidinyl)-4-fluoroaniline, N-(Thiazolyl)-2-(4,6-dimethyl-pyrimidin-2-yl-thio)-acetamide, N-(Thiazolyl)-2-(4-propylphenoxy)-acetamide, 2-[4-(N-Furoyl)aminophenyl]-6-methylbenzothiazole,

- 5 N-(Thiazolyl)-2-(3,5-dimethylphenoxy'-acetamide and 2-[(N-(2-Napthalenyl)amino)-[2,3:5,4]-(5,5-dimethyl-cyclohexanonyl)]thiazole can be purchased from MDD, Inc. (Acton, Ontario), a successor to Ortech Corporation.

In one synthetic process to prepare compounds of the general formula I, a thiazole is reacted with an alkyl or acyl halide in the presence of base such as

- 10 triethylamine, to produce the corresponding alkyl or acyl derivative at the 2 carbon. *See*, Medici et al., *J. Org. Chem.* 49: 590-596, 1984. In some cases, a chromatographic step is applied to separate additions at the 2 and 3 positions of the thiazole ring.


In another synthesis of compounds of the formula I wherein R^c is amino, nitro-containing analogs of compounds of the invention or precursors thereof are

- 15 catalytically hydrogenated to the corresponding amino compounds.

2-Amino thiazole compounds wherein (R^c is amino) can also be synthesized by reacting thiourea (which can be substituted on at least one amine) with an alpha-halo ketone using the method described in *Vogel's Textbook of Practical Organic Chemistry*, 5th Edition, John Wiley & Sons, New York, p. 1153. Such a reaction is exemplified by a

- 20 synthesis of 2-amino-4-phenylthiazole:

Scheme 1




- 25 N-aryl imidazoles can be prepared using appropriate aromatic nucleophilic displacement reactions. For example, fluoro phenyl compounds such as 4-fluorobenzoic acid methyl ester can be used to substitute on the N¹ nitrogen of imidazole to make methyl-4-(1H-imidazol-1-yl)benzoate. *See*, Morgan et al., *J. Med. Chem.* 33: 1091-1097, 1990.

Amino functions of 2-aminoimidazoles or 2-aminothiazoles can be acylated by dehydration or other methods known in the art.


Substituted oxazoles can be prepared by methods known in the art. For instance, 2-unsubstituted oxazoles can be formed by condensation of formamide with either α -hydroxy or α -haloketones intermediates (H. Brederick, R. Gommper, H. G. v. Shuh and G. Theilig, in *Newer Methods of Preparative Organic Chemistry*, Vol. III, ed. W. Foerst, Academic press, New York, 1964, p. 241). The intermediates can cyclize under acid conditions to form the oxazole ring (**Scheme 2**). In addition, 2,4-disubstituted oxazoles can be prepared from α -haloketones and amides at higher temperatures using the same method.

Scheme 2




Oxazoles can be prepared by cyclization reactions of isonitriles (van Leusen, A. M. *Lect. Heterocycl. Chem.* **1980**, *5*, S111; Walborsky, H. M.; Periasamy, M. P. in *The Chemistry of Functional Groups, suppl. C*, Patai, S.; Rappoport, Z., Eds; Wiley-Interscience, 1983, p. 835; Hoppe, D. *Angew. Chem. Int. Edn. Engl.*, **1974**, *13*, 789; Schollkopf, U. *Angew. Chem. Int. Ed. Engl.*, **1977**, *16*, 339). For example, as shown below in **Scheme 3**, the tosylmethyl isocyanide can be deprotonated by a base and reacted with a suitable electrophile (e.g. an aldehyde). The intermediate can cyclize and aromatize to provide the desired oxazole analog. Other methods for preparing oxazoles include 1,5-dipolar cyclization of acylated nitrile ylides (Taylor E. C.; Turchi, I. J. *Chem. Rev.*, **1979**, *79*, 181; Huisgen, R. *Angew. Chem. Int. Edn. Engl.* **1980**, *19*, 947)

Scheme 3




2-Amino-substituted oxazoles (i.e. $\text{R}^c=\text{NH}_2$) can be prepared by two general methods. Urea can be condensed with α -bromo ketones to yield 2-aminooxazoles that can be substituted at the 4 and 5 positions (**Scheme 4**). Alternatively, another route to 2-aminooxazoles from acyclic precursors is the base catalyzed reaction of cyanamide with α -hydroxy ketones (**Scheme 5**). 2-Aminooxazoles of the invention can also be prepared from the nucleophilic displacement of amines with 2-chlorooxazole, for example, for compounds of the invention wherein R^c is ArNH- (Gompper, R.; Effenberger, F. *Chem. Ber.*, **1959**, 92, 1928).

Scheme 4



Scheme 5




15

Compounds of the invention, wherein R^c is arylcarbonyl can be synthesized by acylation of the amino moiety of 2-aminooxazoles with, for example, with anhydrides to yield 2-acylaminoxazoles. In addition, 2-aminooxazoles can be acylated, for instance, with chloroacetic anhydride to yield an α -chloro carboxamide. The α -chloro carboxamide can serve as a suitable alkylating reagent that can be treated with, for example, phenols, arylamines and alkylamines to prepare compounds of the invention. An oxazole of the invention wherein R^c is $\text{Ar-oxycarbonylamino}$ is shown in **Scheme 6**.


20

Scheme 6




Oxazoles of the invention with R^c is aminocarbonylamino (ureido) or aminothiocabonylamino (thioureido) can be prepared from 2-aminooxazoles (**Scheme 7**). 2-Aminooxazoles can be treated with isocyanates and isothiocyanates to yield 2-ureido and 2-thioureido oxazoles, respectively (Crank, G.; Foulis, J. *J. Med. Chem.*, **1971**, *14*, 1075; Crank, G. Neville, M.; Ryden, R. *J. Med. Chem.*, **1974**, *16*, 1402).

Scheme 7




2-Aminooxazoles can be hydrogenated using palladium catalysts to yield 2-aminooxazolines (**Scheme 8**) (Tanaka, C.; Kuriyama, S. *Yakugaku Zasshi* **1979**, *99*, 78).

Scheme 8



Benzoxazole intermediates substituted at the 2 position can be prepared from 2-aminophenols by acylation with, for example, with an acid chloride and cyclization (**Scheme 9**).

Scheme 9



To treat glaucoma or reduced accommodation and associated symptoms, an effective amount of a pharmaceutical compound will be recognized by clinicians but includes an amount effective to treat, reduce, ameliorate, eliminate or prevent one or more symptoms of the disease sought to be treated or the condition sought to be avoided or treated, or to otherwise produce a clinically recognizable change in the pathology of the disease or condition.

In treating glaucoma, agents of the inventions can be administered concurrently or in a combined formulation with one or more α_2 -selective adrenergic agonists, carbonic anhydrase inhibitors or prostaglandin analogs. Examples of α_2 -selective adrenergic agonists include clonidine, apraclonidine, guanfacine, guanabenz and methyldopa, which are administered in effective amounts as is known in the art. Examples of carbonic anhydrase inhibitors include acetazolamide, dichlorphenamide and methazolamide, which are administered in effective amounts as is known in the art. Examples of prostaglandin analogs include PGE₂ and PGF_{2 α} analogs, which are administered in effective amounts as is known in the art, including effective amounts administered by topical application to the eye. Thus, the invention further provides pharmaceutical compositions comprising an agent of the invention in combination with an effective amount of an α_2 -selective adrenergic agonist, carbonic anhydrase inhibitor, prostaglandin analog, or combination thereof.

Pharmaceutical compositions can be prepared to allow a therapeutically effective quantity of the compound of the present invention, and can include a pharmaceutically acceptable carrier, selected from known materials utilized for this purpose. *See, e.g.,* Remington, The Science and Practice of Pharmacy, 1995; Handbook of Pharmaceutical Excipients, 3rd Edition, 1999. Such compositions can be prepared in a variety of forms, depending on the method of administration.

In addition to the subject compound, the compositions of this invention can contain a pharmaceutically-acceptable carrier. The term "pharmaceutically-acceptable carrier", as used herein, means one or more compatible solid or liquid filler diluents or encapsulating substances that are suitable for administration to an animal, including a mammal or human. The term "compatible", as used herein, means that the components of the composition are capable of being commingled with the subject compound, and with each other, such that there is no interaction that would substantially reduce the

pharmaceutical efficacy of the composition under ordinary use. Preferably when liquid dose forms are used, the compounds of the invention are soluble in the components of the composition. Pharmaceutically-acceptable carriers must, of course, be of sufficiently high purity and sufficiently low toxicity to render them suitable for administration to the animal being treated.

Some examples of substances which can serve as pharmaceutically-acceptable carriers or components thereof are sugars, such as lactose, glucose and sucrose; starches, such as corn starch and-potato starch; cellulose and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose, and methyl cellulose; powdered tragacanth; malt; gelatin; talc; solid lubricants, such as stearic acid and magnesium stearate; calcium sulfate; vegetable oils, such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil and oil of theobroma; polyols such as propylene glycol, glycerine, sorbitol, mannitol, and polyethylene glycol; alginic acid; emulsifiers, such as the TweenTM brand emulsifiers; wetting agents, such sodium lauryl sulfate; coloring agents; flavoring agents; tableting agents, stabilizers; antioxidants; preservatives; pyrogen-free water; isotonic saline; and phosphate buffer solutions. The choice of a pharmaceutically-acceptable carrier to be used in conjunction with the subject compound is basically determined by the way the compound is to be administered. If the subject compound is to be injected, the preferred pharmaceutically-acceptable carrier is sterile, physiological saline, with a blood-compatible suspending agent, the pH of which has been adjusted to about 7.4.

If the preferred mode of administering the subject compound is perorally, the preferred unit dosage form is therefore tablets, capsules, lozenges, chewable tablets, and the like. Such unit dosage forms comprise a safe and effective amount of the subject compound, which is preferably from about 0.7 or 3.5 mg to about 280 mg/ 70 kg, more preferably from about 0.5 or 10 mg to about 210 mg/ 70 kg. The pharmaceutically-acceptable carrier suitable for the preparation of unit dosage forms for peroral administration are well-known in the art. Tablets typically comprise conventional pharmaceutically-compatible adjuvants as inert diluents, such as calcium carbonate, sodium carbonate, mannitol, lactose and cellulose; binders such as starch, gelatin and sucrose; disintegrants such as starch, alginic acid and croscarmellose; lubricants such as magnesium stearate, stearic acid and talc. Glidants such as silicon dioxide can be used to improve flow characteristics of the powder-mixture. Coloring agents, such as the FD&C

dyes, can be added for appearance. Sweeteners and flavoring agents, such as aspartame, saccharin, menthol, peppermint, and fruit flavors, are useful adjuvants for chewable tablets. Capsules typically comprise one or more solid diluents disclosed above. The selection of carrier components depends on secondary considerations like taste, cost, and shelf stability, which are not critical for the purposes of this invention, and can be readily made by a person skilled in the art.

Peroral compositions also include liquid solutions, emulsions, suspensions, and the like. The pharmaceutically-acceptable carriers suitable for preparation of such compositions are well known in the art. Such liquid oral compositions preferably comprise from about 0.012% to about 0.933% of the subject compound, more preferably from about 0.033% to about 0.7%. Typical components of carriers for syrups, elixirs, emulsions and suspensions include ethanol, glycerol, propylene glycol, polyethylene glycol, liquid sucrose, sorbitol and water. For a suspension, typical suspending agents include methyl cellulose, sodium carboxymethyl cellulose, cellulose (e.g. Avicel™, RC-591), tragacanth and sodium alginate; typical wetting agents include lecithin and polyethylene oxide sorbitan (e.g. polysorbate 80). Typical preservatives include methyl paraben and sodium benzoate. Peroral liquid compositions may also contain one or more components such as sweeteners, flavoring agents and colorants disclosed above.

Other compositions useful for attaining systemic delivery of the subject compounds include sublingual and buccal dosage forms. Such compositions typically comprise one or more of soluble filler substances such as sucrose, sorbitol and mannitol; and binders such as acacia, microcrystalline cellulose, carboxymethyl cellulose and hydroxypropyl methyl cellulose. Glidants, lubricants, sweeteners, colorants, antioxidants and flavoring agents disclosed above may also be included.

Compositions can also be used to deliver the compound to the site where activity is desired; such as eye drops, gels and creams for ocular disorders.

Compositions of this invention include solutions or emulsions, preferably aqueous solutions or emulsions comprising a safe and effective amount of a subject compound intended for topical intranasal administration. Such compositions preferably comprise from about 0.01% to about 10.0% w/v of a subject compound, more preferably from about 0.1% to about 2.0%. Similar compositions are preferred for systemic delivery of subject compounds by the intranasal route. Compositions intended to deliver the compound systemically by intranasal dosing preferably comprise similar amounts of

a subject compound as are determined to be safe and effective by peroral or parenteral administration. Such compositions used for intranasal dosing also typically include safe and effective amounts of preservatives, such as benzalkonium chloride and thimerosal and the like; chelating agents, such as edetate sodium and others; buffers such as phosphate, citrate and acetate; tonicity agents such as sodium chloride, potassium chloride, glycerin, mannitol and others; antioxidants such as ascorbic acid, acetylcystine, sodium metabisulfite and others; aromatic agents; viscosity adjustors, such as polymers, including cellulose and derivatives thereof; and polyvinyl alcohol and acids and bases to adjust the pH of these aqueous compositions as needed. The compositions may also comprise local anesthetics or other actives. These compositions can be used as sprays, mists, drops, and the like.

Other preferred compositions of this invention include aqueous solutions, suspensions, and dry powders comprising a safe and effective amount of a subject compound intended for atomization and inhalation administration. Such compositions are typically contained in a container with attached atomizing means. Such compositions also typically include propellants such as chlorofluorocarbons 12/11 and 12/114, and more environmentally friendly fluorocarbons, or other nontoxic volatiles; solvents such as water, glycerol and ethanol, these include cosolvents as needed to solvate or suspend the active; stabilizers such as ascorbic acid, sodium metabisulfite; preservatives such as cetylpyridinium chloride and benzalkonium chloride; tonicity adjustors such as sodium chloride; buffers; and flavoring agents such as sodium saccharin. Such compositions are useful for treating respiratory disorders, such as asthma and the like.

Other preferred compositions of this invention include aqueous solutions comprising a safe and effective amount of a subject compound intended for topical intraocular administration. Such compositions preferably comprise from about 0.01% to about 0.8% w/v of a subject compound, more preferably from about 0.05% to about 0.3%. Such compositions also typically include one or more of preservatives, such as benzalkonium chloride or thimerosal; vehicles, such as poloxamers, modified celluloses, povidone and purified water; tonicity adjustors, such as sodium chloride, mannitol and glycerin; buffers such as acetate, citrate, phosphate and borate; antioxidants such as sodium metabisulfite, butylated hydroxy toluene and acetyl cysteine; acids and bases can be used to adjust the pH of these formulations as needed.

Other preferred compositions of this invention useful for peroral administration include solids, such as tablets and capsules, and liquids, such as solutions, suspensions and emulsions (preferably in soft gelatin capsules), comprising a safe and effective amount of a subject compound. Such compositions can be coated by conventional methods, typically with pH or time-dependent coatings, such that the subject compound is released in the gastrointestinal tract at various times to extend the desired action. Such dosage forms typically include, but are not limited to, one or more of cellulose acetate phthalate, polyvinylacetate phthalate, hydroxypropyl methyl cellulose phthalate, ethyl cellulose, EudragitTM coatings, waxes and shellac.

The compounds of the invention are administered by ocular, oral, parenteral, including, for example, using formulations suitable as eye drops. For ocular administration, ointments or droppable liquids may be delivered by ocular delivery systems known to the art such as applicators or eye droppers. Such compositions can include mucomimetics such as hyaluronic acid, chondroitin sulfate, hydroxypropyl methylcellulose or polyvinyl alcohol, preservatives such as sorbic acid, EDTA or benzylchromium chloride, and the usual quantities of diluents and/or carriers. See, Remington's Pharmaceutical Sciences, 16th Ed., Mack Publishing, Easton, PA, 1980, as well as later editions, for information on pharmaceutical compounding.

Numerous additional administration vehicles will be apparent to those of ordinary skill in the art, including without limitation slow release formulations, liposomal formulations and polymeric matrices.

In another preferred embodiment, the pharmaceutically effective amount is approximately 0.1 or 0.5 to 4 mg/kg body weight daily. Still more preferably, the pharmaceutically effective amount is approximately 1 mg/kg body weight daily. In a preferred embodiment, the amount is administered in once daily doses, each dose being approximately 1 mg/kg body weight.

Compounds of the invention can be used in conjunction with monitoring the improvement (decrease) in the intraocular pressure in a mammal using standard methodology.

The methods of the inventions can be assessed in animal models for ophthalmologic function. For example, improvements in fluid outflow facility can be studied in Rhesus monkeys treated with the compounds and methods of the invention. Aged Rhesus monkeys receive a single transcorneal injection of a test compound

(compound of the invention) at a concentration of about 1 mM in the anterior chamber of one eye, and Barany's solution, as a control, in the adjacent eye. Needle outflow facility is measured under baseline and pilocarpine-stimulated conditions at time points (for example, 3, 8, 12 and 24 weeks), after the administration of the test compound.

- 5 Increases in outflow facility in the drug treated vs. the control eye under baseline and cholinergic-stimulated (e.g. pilocarpine) conditions at the various time points are compared. As the enhancement of outflow facility can be influenced by the route of administration of the cholinergic agent, various routes of administration of the cholinergic agent can be used in the experiments. For instance, an intravenous
- 10 administration versus a direct administration of pilocarpine can be compared. The above experiment demonstrates one method of measuring the improvement in ophthalmologic function. Such improvement has been illustrated with 4,5-dimethyl-3-(2-oxoethyl-phenethyl)thiazolium chloride, a compound believed to act by the same mechanism as those described here. See, U.S. application for "Methods for Treating Glaucoma I,"
- 15 concurrently filed herewith.

In addition to measuring increased fluid outflow facility using the methods of the invention, improvements in pilocarpine-stimulated accommodation (i.e, the process of effecting refractive changes in the shape of the lens) can also be assessed in animal studies. As in the regulation of outflow facility, cholinergic input stimulates the

20 movement of the ciliary muscle to control the shape of the lens, and allows accommodation in conditions of low illumination. Accommodation is impaired in a vast majority of individuals and begins to become noticeable to the individual around the age of 40 years. Interestingly, changes in accommodative response occur much earlier in life, around 18 years of age, and progresses until vision is noticeably impaired.

- 25 Physiological studies on accommodation are conducted following intraocular injection of a test compound and the results are compared relative to the results of control (untreated) animals. In the experiment, primates (for example, Rhesus monkeys) are treated twice a day for four days with 2 μ g of prostaglandin F₂ α (PGF₂ α). On days 5-8 both eyes are treated first with 2 μ g of PGF₂ α followed 2 hours later with an
- 30 intraocular injection of 10 μ L of the test compound of a final concentration of 1 mM. No injection is made to the control eye. 24 hours after the last injection of the test compound, a course of therapy consisting of once a day dosing for a total of 4 days

accommodative responses to i.m. pilocarpine administration is performed following phenylephrine refraction. Improvement in accommodation has been illustrated with 4,5-dimethyl-3-(2-oxoethyl-phenethyl)thiazolium chloride, a compound believed to act by the same mechanism as those described here. See, U.S. application for "Methods for Treating Glaucoma I," concurrently filed herewith.

Compounds of the invention can be tested to determine corneal penetration to the anterior chamber of the eye following topical administration of eye drops. For example, a test compound is assayed *in vitro* through an intact rabbit cornea for transcorneal penetration in a standard diffusion chamber apparatus. Corneas are mounted in a chamber at 37 °C with the epithelial side exposed to the test compound in Barany's solution. 1.0 mL samples are taken from the endothelial side 1 hour after addition of the test compound at a final concentration of 1 mM to the epithelial chamber. The volume of the chamber is replaced with phosphate buffered saline. The amount of test compound can be measured using any means that can be used to separate the compound and measure its concentration. For example, an HPLC with an attached UV detector can be used to determine the concentration of the test compound that has penetrated the cornea. Penetration values are also determined at later time points, for example, at 5 hours.

Assessment of corneal penetration of compounds of the invention can be determined *in vivo*, for example, in Cynomolgus monkeys. During these studies, the penetration of a test compound is evaluated using an eye-cup which holds a solution of 10 mM of the test compound in Barany's solution for 5 hours. At the end of the experiment the eye cup is removed, the eye is repeatedly flooded with Barany's solution and a sample of intraocular fluid is removed from the anterior chamber with a needle inserted through the cornea. The quantity of the test compound in the intraocular fluid is determined using, for example, HPLC methods.

The activity of the compounds of the invention in breaking, reversing or inhibiting the formation of AGE's or AGE-mediated cross-links can be assayed by any of the methods described in US Patent 5,853,703.

The following examples further illustrate the present invention, but of course, should not be construed as in any way limiting its scope.

EXAMPLE 1. 2,6-diamino-benzothiazole dihydrochloride: 4 g of 2-amino-6-nitrobenzothiazole (Aldrich) was suspended in 130 ml MeOH, and 0.4 g 10% Pd/c (Aldrich) added. The suspension was hydrogenated at room temperature under 60 psi H₂ for 6.5 h. The reaction mixture was filtered, and the particulate washed with MeOH.

- 5 The filtrate was concentrated under reduced pressure, and crystals formed from the concentrate were collected to yield 2.67 g. mp 196-198°C, yield 81.6%. 0.91 g of this product was dissolved in 22 ml MeOH, and the pH adjusted with HCl to 4 to produce 1.2 g of crystals of 2,6-diamino-benzothiazole dihydrochloride. mp 318-320°C, 92.3% yield. Anal. calc. for C₇H₉N₃SCl₂, C 35.30%, H 3.80%, N 17.64%. Found, C 34.91%, H 3.67%, N 17.71%.
- 10

EXAMPLE 2. 2-(3,5-Dimethylphenoxy)-N-thiazol-2-yl)acetamide: First Route: 3,5-Dimethylphenol is reacted with bromoacetic acid at 110°C for four hours, with the reaction mixture stirred overnight without added heat. The resulting (3,5-

15 dimethylphenoxy)acetic acid is dissolved in methylene chloride and coupled to 2-aminothiazole in an overnight, room temperature reaction conducted in the presence of base (N-methylmorpholine) and dehydration mediators 1-hydroxybenzotriazole and 1-(3-dimethylaminopropyl)-3-ethyl-carbodiimide hydrochloride.

- Second Route: 3,5-Dimethylphenol is reacted for 4.5 h with bromoacetic acid in
- 20 THF under nitrogen and in the presence of sodium hydride. The resulting (3,5-dimethylphenoxy)acetic acid is reacted overnight with thionyl chloride, with heat. The resulting (3,5-dimethylphenoxy)acetyl chloride is reacted overnight with 2-aminothiazole in the presence of triethylamine, with cooling to 0°C.

- Third Route: 2-Aminothiazole (20g, 199.7 mmol) was suspended in methylene
- 25 chloride (200 ml), in the presence of pyridine (20 ml, ~250 mmol), and the mixture cooled to 0°C. Bromoacetyl bromide (18.1 ml, 207.6 mmol) was dissolved in 400 ml methylene chloride, and this solution added to the suspended 2-aminothiazole dropwise. The resulting reaction mixture was stirred at room temperature overnight. The crude product was washed with water (200 ml, 1X), then sodium bicarbonate solution (200 ml,
- 30 2X), dried over Na₂SO₄, filtered, and evaporated. The product 2-bromoacetamidothiazole was crystallized from MeOH. Yield, 4g. mp 148°C.

A solution of 3,5-dimethylphenol (2.5g, 13.9 mmol) in dry DMF (20 ml) was placed under a dry nitrogen atmosphere. Sodium hydride (0.7 g, 27.8 mmol; a 60% dispersion in mineral oil) was added in portions, and the mixture stirred for 1 h. A solution of 2-bromoacetamidothiazole (3.0 g, 13.9 mmol) in dry DMF (10 ml) was added to the mixture dropwise. The reaction was heated to 90°C for 5 h, then maintained overnight without external heat. The reaction mixture was poured into ice water, and the resulting material extracted with methylene chloride (50 ml X 3). The organic layer was washed with water (100 ml X 5), dried over Na₂SO₄, filtered, and evaporated. The residue from evaporation was purified by silica gel chromatography developed with pet. ether: ether (1:1 v/v). The product N-(Thiazolyl)-2-(3,5-dimethylphenoxy)-acetamide was crystallized from acetonitrile and methyl tert-butyl ether. Yield, 1.04 g. mp 124-125 °C.

EXAMPLE 3. 2-Furyl-N-[4-(6-methyl-benzothiazol-2-yl)phenyl]carboxamide: First Route: 2-Furoic acid (1.85 gms, 16.5 mmole) was dissolved in anhydrous methylene chloride (30 mls), to which solution was added a suspension of 2-(4-amino-phenyl)-6-methyl benzothiazole (4.76 g., 16.5 mmole) and N-methyl morpholine (2.0 g., 16.5 mmole) in methylene chloride (30 mls, at room temperature). Then, 1-hydroxy-benzotriazole hydrate (2.67 g., 16.5 mmole) and 1-(3-dimethyl amino propyl)-3-ethyl carbodiimide hydrochloride (4.75 g., 16.5 mmole) were added at room temperature. More methylene chloride (20 ml.) was added with stirring at room temperature, and the reaction maintained overnight. The initial clear reaction solution changed to a turbid solution. More methylene chloride (10 ml.) was added to the product mixture, which was then extracted with 1N HCl to separate a solid. The solid was filtered and washed with water. The product solid was crystallized from large amount of MeOH to yield 2.19 gm. (33.1%). mp. 238-240°C. ¹H and ¹³C NMR were consistent with the expected product. TLC showed one spot (5% MeOH-CH₂Cl₂ as developing solvent on silica gel plate).

Route 2: 2-(4-aminophenyl)-6-methyl benzothiazole (2.0 gm, 8.3 mmole) and 2-Furoyl chloride (1.086 gm., 8.32 mmole) were suspended in methylene chloride (30 ml, anhydrous). triethylamine (1.24 gm., 12.25 mmole) was added to the reaction mixture with stirring at room temperature for 2 days. (pH 7.0-7.2). Methylene chloride (50 ml)

was added to the reaction mixture, and the reaction mixture extracted with 1 N HCl (50 ml) to separate a solid. The solid was filtered and washed with water to yield 1.3 gm. (46%) of the desired compound. The product was crystallized from MeOH to obtain 0.99 gm. mp. 238-240°C. ¹H and ¹³C NMR were consistent with the expected product.

5 TLC showed one spot (5% MeOH-CH₂Cl₂ as developing solvent on silica gel plate).

EXAMPLE 4. Cross-Linking Inhibition Assay

The following method was used to evaluate the ability of the compounds to inhibit the cross-linking of glycated bovine serum albumin (AGE-BSA) to rat tail tendon collagen-coated 96-well plates.

10

AGE-BSA was prepared by incubating BSA at a concentration of 200 mg per ml with 200 mM glucose in 0.4M sodium phosphate buffer, pH 7.4 at 37°C for 12 weeks. The glycated BSA was then extensively dialyzed against phosphate buffer solution (PBS) for 48 hours with additional 5 times buffer exchanges. The rat tail tendon collagen coated plate was blocked first with 300 microliters of Superbloc blocking buffer (Pierce Chemical, Rockford, IL) for one hour. The blocking solution was removed from the wells by washing the plate twice with phosphate buffered saline (PBS)-Tween 20 solution (0.05% Tween 20) using a NUNC-multiprobe (Nalge Nunc, Rochester, NY) or Dynatech ELISA-plate (Dynatech, Alexandria, VA) washer. Cross-linking of AGE-BSA (1 to 10 microgram per well depending on the batch of AGE-BSA) to rat tail tendon collagen coated plate was performed with and without the testing compound dissolved in PBS buffer at pH 7.4 at one or more desired concentrations by the addition of 50 microliters each of the AGE-BSA diluted in PBS or in the solution of test compound at 37°C for 4 hours. Unbrowned BSA in PBS buffer with or without testing compound were added to the separate wells as the blanks. The un-cross-linked AGE-BSA was then removed by washing the wells three times with PBS-Tween buffer. The amount of AGE-BSA crosslinked to the tail tendon collagen-coated plate was then quantitated using a polyclonal antibody raised against AGE-RNase. After a one-hour incubation period, AGE antibody was removed by washing 4 times with PBS-Tween.

15
20
25
30

The bound AGE antibody was then detected with the addition of horseradish peroxidase-conjugated secondary antibody—goat anti-rabbit immunoglobulin and incubation for 30 minutes. The substrate of 2,2-azino-di(3-ethylbenzthiazoline sulfonic

acid) (ABTS chromogen) (Zymed Laboratories, Inc., South San Francisco, CA) was added. The reaction was allowed for an additional 15 minutes and the absorbance was read at 410 nm in a Dynatech plate reader.

5 EXAMPLE 5 Cross-Link Breaking Assay

To ascertain the ability of the compounds of the instant invention to break or reverse already formed advanced glycosylation endproducts, a sandwich enzyme immunoassay was applied. Generally, the assay utilizes collagen-coated 96 well microtiter plates that are obtained commercially. AGE-modified protein (AGE-BSA) is incubated on the collagen-coated wells for four hours, is washed off the wells with PBS-Tween and solutions of the test compounds are added. Following an incubation period of 16 hours (37°C) cross-link-breaking is detected using an antibody raised against AGE-ribonuclease or with an antibody against BSA.

Preparation of solutions and buffers

15 Bovine Serum Albumin (Type V) (BSA) (from Calbiochem) solution was prepared as follows: 400 mg of Type V BSA (bovine serum albumin) was added for each ml of 0.4 M sodium phosphate buffer, pH 7.4. A 400 mM glucose solution was prepared by dissolving 7.2 grams of dextrose in 100 ml of 0.4 M sodium phosphate buffer, pH 7.4. The BSA and glucose solutions were mixed 1:1 and incubated at 37°C for 12 weeks.

20 The pH of the incubation mixture was monitored weekly and adjusted to pH 7.4 if necessary. After 12 weeks, the AGE-BSA solution was dialyzed against PBS for 48 hours with four buffer changes, each at a 1:500 ratio of solution to dialysis buffer. Protein concentration was determined by the micro-Lowry method. The AGE-BSA stock solution was aliquoted and stored at -20°C.

25 Test compounds were dissolved in PBS and the pH was adjusted to pH 7.4, if necessary. AGE-BSA stock solution was diluted in PBS to measure maximum crosslinking and in the inhibitor solution for testing inhibitory activity of compounds. The concentration of AGE-BSA necessary to achieve the optimum sensitivity was determined by initial titration of each lot of AGE-BSA.

30 Substrates for detection of secondary antibody binding were prepared by diluting the HRP substrate buffer (Zymed) 1:10 in distilled water and mixing with ABTS chromogen (Zymed) 1:50 just prior to use.

Assay Procedures

Biocoat plates were blocked with 300 microliters of Superbloc (Pierce Chemical). Plates were blocked for one hour at room temperature and were washed with PBS-Tween (0.05% v/v) three times with the Dynatech platewasher before addition of test reagents.

5 The first three wells of the Biocoat plate were used for the reagent blank. Fifty microliters of solutions AGE-BSA were added to test wells in triplicate and only PBS in blank wells. The plate was incubated at 37°C for four hours and washed with PBS-Tween three times. Fifty microliters of PBS was added to the control wells and 50 microliters of the test prospective agent was added to the test wells and blank. The plate
10 was incubated overnight (approximately 16 hours) with prospective agent, followed by washing in PBS before addition of primary antibody.

(Prior to use, each lot of primary antibody, either anti-BSA or anti-RNase, was tested for optimum binding capacity in this assay by preparing serial dilutions (1:500 to 1:2000) and plating 50 microliters of each dilution in the wells of Biocoat plates.

15 Optimum primary antibody was determined from saturation kinetics.) Fifty microliters of primary antibody of appropriate dilution, was added and incubated for one hour at room temperature. The plate was then washed with PBS-Tween.

Plates were incubated with the secondary antibody, HRP-(Goat-anti-rabbit), which was diluted 1:4000 in PBS and used as the final secondary antibody. The
20 incubation was performed at room temperature for thirty minutes.

Detection of maximum crosslinking and breaking of AGE crosslinking was performed as follows. HRP substrate (100 microliter) was added to each well of the plate and was incubated at 37°C for fifteen minutes. Readings were taken in the Dynatech ELISA-plate reader.

25

Definitions

Heterocycle. Except where heteroaryl is separately recited for the same substituent, the term "heterocycle" includes heteroaryl.

All publications and references, including but not limited to patents and patent
30 applications, cited in this specification are herein incorporated by reference in their entirety as if each individual publication or reference were specifically and individually indicated to be incorporated by reference herein as being fully set forth. Any patent

application to which this application claims priority is also incorporated by reference herein in its entirety in the manner described above for publications and references.

While this invention has been described with an emphasis upon preferred embodiments, it will be obvious to those of ordinary skill in the art that variations in the preferred devices and methods may be used and that it is intended that the invention may
5 be practiced otherwise than as specifically described herein. Accordingly, this invention includes all modifications encompassed within the spirit and scope of the invention as defined by the claims that follow.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
22